

## RESOURCES

- Download this card and additional resources at [www.sprc.org](http://www.sprc.org) or at [www.stopasuicide.org](http://www.stopasuicide.org)
- Resource for implementing The Joint Commission 2007 Patient Safety Goals on Suicide [www.sprc.org/library/jcsafetygoals.pdf](http://www.sprc.org/library/jcsafetygoals.pdf)
- SAFE-T drew upon the American Psychiatric Association Practice Guidelines for the Assessment and Treatment of Patients with Suicidal Behaviors [www.psychiatryonline.com/pracGuide/pracGuideTopic\\_14.aspx](http://www.psychiatryonline.com/pracGuide/pracGuideTopic_14.aspx)
- Practice Parameter for the Assessment and Treatment of Children and Adolescents with Suicidal Behavior. Journal of the American Academy of Child and Adolescent Psychiatry, 2001, 40 (7 Supplement): 24s-51s

## ACKNOWLEDGEMENTS

- Originally conceived by Douglas Jacobs, MD, and developed as a collaboration between Screening for Mental Health, Inc. and the Suicide Prevention Resource Center.
- This material is based upon work supported by the Substance Abuse and Mental Health Services Administration (SAMHSA) under Grant No. 1U79SM57392. Any opinions/ findings/conclusions/recommendations expressed in this material are those of the author and do not necessarily reflect the views of SAMHSA.

National Suicide Prevention Lifeline  
**1.800.273.TALK (8255)**

COPYRIGHT 2009 BY EDUCATION DEVELOPMENT CENTER, INC. AND  
SCREENING FOR MENTAL HEALTH, INC. ALL RIGHTS RESERVED.  
PRINTED IN THE UNITED STATES OF AMERICA.  
FOR NON-COMMERCIAL USE.



[www.sprc.org](http://www.sprc.org)



[www.mentalhealthscreening.org](http://www.mentalhealthscreening.org)

# SAFE-T

## Suicide Assessment Five-step Evaluation and Triage

*for Mental Health Professionals*

1

IDENTIFY RISK FACTORS  
Note those that can be  
modified to reduce risk

2

IDENTIFY PROTECTIVE FACTORS  
Note those that can be enhanced

3

CONDUCT SUICIDE INQUIRY  
Suicidal thoughts, plans  
behavior and intent

4

DETERMINE RISK LEVEL/INTERVENTION  
Determine risk. Choose appropriate  
intervention to address and reduce risk

5

DOCUMENT  
Assessment of risk, rationale,  
intervention and follow-up

NATIONAL SUICIDE PREVENTION LIFELINE  
**1.800.273.TALK (8255)**

*Suicide assessments should be conducted at first contact, with any subsequent suicidal behavior, increased ideation, or pertinent clinical change; for inpatients, prior to increasing privileges and at discharge.*

## 1. RISK FACTORS

- ✓ **Suicidal behavior:** history of prior suicide attempts, aborted suicide attempts or self-injurious behavior
- ✓ **Current/past psychiatric disorders:** especially mood disorders, psychotic disorders, alcohol/substance abuse, ADHD, TBI, PTSD, Cluster B personality disorders, conduct disorders (antisocial behavior, aggression, impulsivity).  
*Co-morbidity and recent onset of illness increase risk*
- ✓ **Key symptoms:** anhedonia, impulsivity, hopelessness, anxiety/panic, insomnia, command hallucinations
- ✓ **Family history:** of suicide, attempts or Axis 1 psychiatric disorders requiring hospitalization
- ✓ **Precipitants/Stressors/Interpersonal:** triggering events leading to humiliation, shame or despair (e.g., loss of relationship, financial or health status—real or anticipated). Ongoing medical illness (esp. CNS disorders, pain). Intoxication. Family turmoil/chaos. History of physical or sexual abuse. Social isolation.
- ✓ **Change in treatment:** discharge from psychiatric hospital, provider or treatment change
- ✓ **Access to firearms**

## 2. PROTECTIVE FACTORS *Protective factors, even if present, may not counteract significant acute risk*

- ✓ **Internal:** ability to cope with stress, religious beliefs, frustration tolerance
- ✓ **External:** responsibility to children or beloved pets, positive therapeutic relationships, social supports

## 3. SUICIDE INQUIRY *Specific questioning about thoughts, plans, behaviors, intent*

- ✓ **Ideation:** frequency, intensity, duration--in last 48 hours, past month and worst ever
- ✓ **Plan:** timing, location, lethality, availability, preparatory acts
- ✓ **Behaviors:** past attempts, aborted attempts, rehearsals (tying noose, loading gun), vs. non-suicidal self injurious actions
- ✓ **Intent:** extent to which the patient (1) expects to carry out the plan and (2) believes the plan/act to be lethal vs. self-injurious; Explore ambivalence: reasons to die vs. reasons to live

*\* For Youths: ask parent/guardian about evidence of suicidal thoughts, plans, or behaviors, and changes in mood, behaviors or disposition*

*\* Homicide Inquiry: when indicated, esp. in character disordered or paranoid males dealing with loss or humiliation. Inquire in four areas listed above.*

## 4. RISK LEVEL/INTERVENTION

- ✓ **Assessment** of risk level is based on clinical judgment, after completing steps 1-3
- ✓ **Reassess** as patient or environmental circumstances change

RISK LEVEL	RISK / PROTECTIVE FACTOR	SUICIDALITY	POSSIBLE INTERVENTIONS
High	Psychiatric disorders with severe symptoms, or acute precipitating event; protective factors not relevant	Potentially lethal suicide attempt or persistent ideation with strong intent or suicide rehearsal	Admission generally indicated unless a significant change reduces risk. Suicide precautions
Moderate	Multiple risk factors, few protective factors	Suicidal ideation with plan, but no intent or behavior	Admission may be necessary depending on risk factors. Develop crisis plan. Give emergency/crisis numbers
Low	Modifiable risk factors, strong protective factors	Thoughts of death, no plan, intent or behavior	Outpatient referral, symptom reduction. Give emergency/crisis numbers

(This chart is intended to represent a range of risk levels and interventions, not actual determinations.)

## 5. DOCUMENT Risk level and rationale; treatment plan to address/reduce current risk (e.g., setting, medication, psychotherapy, E.C.T., contact with significant others, consultation); firearm instructions, if relevant; follow up plan. For youths, treatment plan should include roles for parent/guardian.